

Vaccine	Min. # of vaccine doses by childcare and pre-school age levels							Vaccine doses by school grade level												Notes			
	By 4 Mo.	By 6 Mo.	By 12 Mo.	By 15 Mo.	16-47 Mo.	48-59 Mo.	≥ 60 Mo.	K	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th	8 th	9 th	10 th	11 th		12 th		
Diphtheria/Tetanus/Pertussis (DTP/DTaP/TD)*	1	2	3	3	3	4	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	1 dose required on/after 4th birthday. 4 doses sufficient if last dose given on/after 4th birthday.
Tetanus/Diphtheria/Pertussis (Tdap)																1	1	1	1	1	1	1	1 dose Tdap required for entry into 7th grade, Meningococcal (ACWY) and HPV vaccinations are also recommended at the same time.
Polio (OPV/IPV)*	1	2	3	3	3	4 (3)	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	3 doses sufficient if last dose was given on/after 4th birthday. Students in K through 7th grades final dose required on or after 4th birthday.
Measles/Mumps/Rubella (MMR)				1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	Minimum age for valid 1st dose is 12 months. Live vaccines (MMR, Varicella) must be given on the same day; if not, they must be administered a minimum of 28 days apart.
Haemophilus Influenzae type B (Hib)*	1	2	2	2	3/2/1	3/2/1	3/2/1																
Hepatitis B (HepB)	1	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	Two doses adult Recombivax HB is also valid if administered at ages 11-15 yrs. and if 2nd dose received no sooner than 16 weeks after 1st dose.
Pneumococcal (PCV)*	2	3	3	4/3/2/1	4/3/2/1	4/3/2/1	4/3/2/1/0																
Varicella (VAR)**				1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1	1	Min. age for 1st dose 12 months; 2 doses recommended for 11th-12th graders. 2nd dose should ideally be given after 4th b-day, see note below . Live vaccines (MMR, Varicella) must be given on the same day; if not, they must be administered a min. of 28 days apart. For K-2nd graders, receipt of vaccine, titer or laboratory-confirmed diagnosis is required as proof of prior disease.
Hepatitis A (HepA)				1	1	1	1																

Diphtheria/Tetanus/Pertussis: If child (4 months-6 years) is behind schedule, follow the CDC's catchup schedule.

Tetanus/Diphtheria/Pertussis: Students in 7th-12th grades require proof of 1 dose Tdap received regardless of when the last Td-containing vaccine was given.

Children 7-18 yrs. who are not fully immunized with the childhood DTaP series should be vaccinated according to the CDC's catch-up schedule, with Tdap preferred as the first dose followed by Td if needed. A 3-dose series is sufficient if initiated after 7 yrs. of age in which 1 dose must be Tdap, followed by 2 doses of Td.

Polio: A minimum of 4 weeks between doses required with 6 months between last 2 doses.

MMR: Required second dose should be given on/after 4th birthday. However, dose 2 may be given earlier with at least 4 weeks between 1st and 2nd doses.

Hib: If series started <12 months of age, 3 doses required with at least 1 dose on/after 1st birthday. 2 doses required if 1st dose received at 12-14 months. 1 dose of Hib vaccine administered between 16 and 59 months of age is sufficient. Not recommended ≥60 months of age.

Hep B: 2nd dose a minimum of 4 weeks after dose 1; 3rd dose at least 16 weeks after dose 1 and at least 8 weeks after dose 2. Infants currently receiving primary series, final dose should be administered no earlier than age 24 weeks.

PCV: Administer a series of PCV13 vaccine at ages 2, 4, 6 months with a booster at age 12-15 months. Catch-up: Administer 1 dose of PCV13 to all healthy children aged 12-59 months who are not completely vaccinated for their age; children >60 months, no doses required.

Varicella: For children aged 12 months-12 years, the minimum interval between the two doses is 3 months. However, if dose 2 was administered ≥28 days after dose 1, dose 2 is considered valid and need not be repeated. For children ≥13 years, the recommended minimum interval is 4 wks. Required for proof of varicella immunity: Receipt of vaccine, written proof of immunity by a physician/health care provider or laboratory titer.

For K through 2nd graders, receipt of vaccine, titer or laboratory confirmed diagnosis is required as proof of prior disease. For all newly diagnosed Varicella cases laboratory confirmation of disease is required.

Hep A: One dose required by 15 months; two doses recommended with at least 6 months between doses. Hepatitis A vaccine is only required for childcare and pre-school age children, but is recommended for all ages.

Recommended schedule for adult students 19+ years of age in secondary school	
Vaccine	Recommended doses
Tetanus/Diphtheria/Pertussis (TdaP)	1
Measles/Mumps/Rubella (MMR)	2
Varicella (VAR)**	2

Resources
NM Immunization Protocol: https://nmhealth.org/publication/view/regulation/531/
NMSIIS: https://nmsiis.health.state.nm.us
NM School Health Manual: http://www.nmschoolhealthmanual.org
<i>For more information call your Regional DOH Immunization contact</i>

*Minimum age 6 weeks. Note: Any vaccine administered ≤4 days prior to minimum interval or age is valid. All students enrolled in designated grades are expected to meet requirements.

**In the event of an outbreak two doses may be required for all students; students who have had only one dose may be excluded from school.